International Journal of Scientific Research and Modern Technology (IJSRMT) ijsrmt.com
Volume 1, Issue 11, 2022
DOI: https://doi.org/10.38124/ijsrmt.v1i11.1785

Quantitative Assessment of Laboratory Documentation Errors and Training Effectiveness in Strengthening Good Manufacturing Practice Compliance

Himabindhu Kondeti[1]

[1]Department of Quality Assurance, Independent Researcher, Actalent., St. Louis, Missouri, USA

Publication Date: 2022/11/28

Abstract

Laboratory documentation errors constitute a significant source of Good Manufacturing Practice (GMP) non-compliance, compromising data integrity, analytical traceability, product quality assurance, and regulatory inspection readiness in pharmaceutical manufacturing environments. Conventional documentation control systems frequently depend on retrospective audits, manual error classification, periodic personnel training, and descriptive compliance assessments, which provide limited capacity for predicting documentation failures and quantitatively evaluating training effectiveness. This study proposes a novel Documentation Error Risk and Training Optimization Algorithm (DERTOA), integrating machine learning, statistical process control, personnel competency assessment, and multi-objective optimization to quantify documentation errors, predict non-compliance risks, and optimize training interventions in GMP-regulated laboratories. The proposed framework evaluates documentation error frequency, error severity, correction timeliness, training completion, competency retention, procedural adherence, and audit non-conformity rates using laboratory quality assurance records, training histories, and inspection observations. DERTOA combines an error-severity-weighted compliance risk model with gradient-boosted classification and adaptive training allocation to identify personnel, documentation activities, and laboratory operations requiring targeted corrective interventions. Its predictive and optimization performance is designed to be evaluated against Logistic Regression (LR), Random Forest (RF), Extreme Gradient Boosting (XGBoost), and conventional rule-based training allocation, using accuracy, precision, recall, F1-score, area under the receiver operating characteristic curve (AUC-ROC), mean absolute error (MAE), and root mean squared error (RMSE), where appropriate. A longitudinal quantitative evaluation incorporating pre-training and post-training observations, interrupted time-series analysis, and controlled simulation experiments is proposed to estimate documentation error reduction, competency improvement, training resource utilization, and sustained GMP compliance. Comparative performance graphs, error-distribution histograms, compliance trend charts, confusion matrices, and training effectiveness heatmaps will characterize differences between DERTOA and benchmark approaches. The anticipated contribution is an interpretable, risk-adaptive decision-support framework that links documentation error prevention with competency-based training and measurable compliance improvement. By integrating predictive analytics with targeted training optimization, the proposed methodology aims to strengthen laboratory data integrity, reduce recurrent documentation deficiencies, and support continuous GMP compliance monitoring in pharmaceutical quality control laboratories.

Keywords: *Laboratory Documentation Errors; Training Effectiveness; Good Manufacturing Practice; Compliance Assessment; Data Integrity.*

I. INTRODUCTION

- *Background of Good Manufacturing Practice Compliance, Laboratory Documentation Systems, and Pharmaceutical Quality Assurance Requirements*

Good Manufacturing Practice (GMP) compliance constitutes a fundamental regulatory requirement for ensuring pharmaceutical products consistently satisfy established standards of identity, strength, purity, quality, and safety. Within pharmaceutical quality control laboratories, compliance depends on validated analytical procedures, qualified equipment, controlled documentation, competent personnel, and reliable quality management systems. Laboratory documentation provides

Kondeti, H. (2022). Quantitative Assessment of Laboratory Documentation Errors and Training Effectiveness in Strengthening Good Manufacturing Practice Compliance. *International Journal of Scientific Research and Modern Technology*, *1*(11), 57–77. https://doi.org/10.38124/ijsrmt.v1i11.1785

the traceable evidence necessary to demonstrate that analytical activities have been performed according to approved procedures and established specifications. These records include analytical worksheets, laboratory notebooks, instrument calibration histories, chromatographic data, sample preparation records, deviation investigations, and electronic audit trails. Yu et al. (2014) established that systematic pharmaceutical quality management requires scientifically justified control strategies, identification of critical quality attributes, and continuous evaluation of manufacturing processes. Accordingly, laboratory documentation must satisfy the principles of attributable, legible, contemporaneous, original, and accurate records, supplemented by completeness, consistency, endurance, and availability. Documentation failures can compromise analytical reproducibility, batch disposition decisions, and regulatory confidence even when the underlying experimental procedure has been performed correctly.

Modern pharmaceutical manufacturing increasingly employs laboratory information management systems, electronic laboratory notebooks, computerized analytical platforms, and integrated quality management software to strengthen operational traceability and compliance oversight. Nevertheless, technological automation does not independently eliminate transcription errors, delayed entries, incomplete records, unauthorized modifications, or procedural deviations. Rantanen and Khinast (2015) emphasized the growing importance of advanced manufacturing technologies, process understanding, and data-driven quality assurance within pharmaceutical production systems. Extending this perspective, the present study conceptualizes GMP documentation compliance as a measurable interaction among record accuracy, personnel competency, analytical process reliability, and organizational quality controls. The proposed Documentation Error Risk and Training Optimization Algorithm (DERTOA) integrates severity-weighted documentation error assessment with predictive analytics and competency-based training allocation. Its framework will quantify documentation deficiencies, evaluate their regulatory implications, and identify opportunities for targeted corrective interventions. This approach establishes the methodological basis for comparing conventional compliance monitoring against integrated predictive and training optimization strategies using documentation error rates, compliance indices, classification performance, and longitudinal training effectiveness indicators.

- *Prevalence, Classification, Severity, and Operational Consequences of Laboratory Documentation Errors in GMP-Regulated Manufacturing Environments*

Laboratory documentation errors represent persistent quality system deficiencies that can undermine the scientific validity, traceability, and regulatory acceptability of pharmaceutical analytical records. These errors encompass missing analytical entries, incorrect sample identification, transcription inaccuracies, delayed result recording, incomplete calculations, unauthorized data alteration, inadequate audit trails, and improper documentation corrections. Their prevalence varies according to laboratory workload, documentation complexity, personnel experience, computerized system configuration, and management oversight. Rogers et al. (2020) examined 300 warning letters issued through the United States Food and Drug Administration's Bioresearch Monitoring Program and identified inadequate documentation and failure to follow established procedures among recurring regulatory deficiencies. Although the investigation principally concerned premarket regulatory activities rather than routine GMP manufacturing laboratories, its findings demonstrate the broader regulatory significance of incomplete and unreliable records. For pharmaceutical quality control operations, documentation errors should therefore be classified according to occurrence frequency, affected analytical process, regulatory criticality, recurrence, detectability, and potential consequences for product quality decisions.

The operational consequences of documentation deficiencies extend beyond administrative non-conformance to analytical result invalidation, delayed batch release, repeated testing, corrective and preventive action investigations, increased inspection exposure, and potential product quality risks. Jaiswal et al. (2020) identified data integrity vulnerabilities associated with backdating, duplicated records, inadequate procedures, compromised electronic access controls, and questionable analytical data management practices. Such deficiencies may exhibit different levels of severity: minor errors involve isolated procedural discrepancies without demonstrable effects on data reliability; major errors compromise documentation completeness or traceability; and critical errors threaten data authenticity, product quality evaluation, or regulatory decision-making. These classifications must be confirmed through documented risk assessment rather than predetermined solely by error type. The present study proposes a standardized severity-weighted documentation error index combining normalized occurrence rates, consequence scores, recurrence indicators, and correction delays. Laboratory activities including sample registration, weighing, chromatographic analysis, result calculation, and analytical review will be evaluated comparatively. Error-distribution histograms, Pareto charts, and severity heatmaps will identify high-risk documentation processes, while longitudinal analyses will assess whether targeted training reduces recurrent deficiencies and improves compliance performance relative to conventional documentation control approaches.

- *Personnel Training Effectiveness, Competency Retention, Documentation Accuracy, and Regulatory Compliance Performance*

Personnel training effectiveness constitutes a critical determinant of laboratory documentation accuracy, procedural consistency, and sustainable GMP compliance because analytical reliability depends substantially on employees' ability to interpret, execute, and document

approved laboratory procedures correctly. Conventional pharmaceutical training programs commonly incorporate initial GMP induction, standard operating procedure instruction, annual refresher courses, and competency assessments following procedural modifications. However, training completion alone does not establish whether acquired knowledge translates into consistent documentation behavior under operational conditions. Hughes et al. (2018) emphasized that organizational training achieves practical value when acquired competencies are transferred into observable workplace performance. Within pharmaceutical laboratories, such transfer involves accurate sample identification, contemporaneous analytical recording, appropriate data correction, complete investigation documentation, and proper electronic record management. Consequently, training effectiveness should be evaluated through multiple indicators, including pre-training and post-training competency scores, practical assessment performance, documentation error rates, corrective action recurrence, and time-dependent knowledge retention. These measurements enable differentiation between theoretical knowledge acquisition and sustained improvements in regulatory compliance behavior.

Training effectiveness also depends on workplace supervision, organizational support, personnel motivation, workload distribution, and opportunities to reinforce acquired competencies. Hughes et al. (2020) demonstrated through meta-analysis that peer support, supervisory support, and organizational support contribute positively to training transfer and sustained workplace application. Their findings support the incorporation of organizational context into documentation competency assessment rather than treating training attendance as an isolated compliance indicator. Building upon this evidence, DERTOA is designed to integrate individual competency profiles, historical documentation error patterns, task criticality, and knowledge retention indicators within an adaptive training optimization framework. Training requirements will be prioritized according to predicted documentation risk and identified competency deficiencies while considering available training resources. Effectiveness will be evaluated using repeated competency assessments, error rates per standardized documentation opportunity, and pre-intervention versus post-intervention compliance measurements. The proposed analysis will distinguish immediate learning gains from persistent behavioral improvement and employ statistical comparisons to determine whether personalized training allocation offers measurable advantages over standardized refresher schedules in reducing documentation deficiencies across laboratory operations.

- *Limitations of Conventional Documentation Audits, Rule-Based Error Detection, and Standardized GMP Training Evaluation Approaches*

Conventional laboratory documentation compliance assessments predominantly rely on scheduled internal audits, manual record verification, checklist-based inspections, deviation reporting, and retrospective corrective action investigations. Although these approaches remain essential regulatory controls, their effectiveness may be constrained by inspection sampling, delayed error identification, auditor subjectivity, inconsistent classification criteria, and limited integration of historical compliance information. West et al. (2017) identified methodological inconsistencies in laboratory error recording and emphasized the importance of standardized quality indicators, systematic data capture, and statistical process monitoring. Their review concerned medical laboratory preanalytical processes; nevertheless, the methodological observations are relevant to developing structured documentation error surveillance in pharmaceutical laboratories. Traditional rule-based systems typically identify predefined deficiencies, such as missing signatures, incomplete mandatory fields, expired training records, or deviations from specified procedural thresholds. Such systems may provide reliable detection of explicitly encoded violations but struggle to recognize interacting risk factors, recurrent behavioral patterns, and emerging documentation vulnerabilities that are not represented within established decision rules.

A further limitation concerns the separation of documentation error monitoring from training effectiveness evaluation. Conventional training systems frequently emphasize attendance, examination completion, and scheduled refresher participation without directly linking individual competency development to subsequent documentation performance. Schmitt et al. (2020) demonstrated the potential of predictive machine learning approaches to support targeted quality inspection and improve operational decision-making in industrial manufacturing. Although their application was not developed specifically for pharmaceutical documentation compliance, the underlying predictive inspection concept provides a methodological basis for investigating risk-adaptive laboratory oversight. The proposed DERTOA framework addresses these limitations by combining severity-weighted error classification, nonlinear risk prediction, temporal competency assessment, and constrained training optimization. Comparative experiments will evaluate Logistic Regression, Random Forest, XGBoost, and conventional rule-based training allocation against the proposed integrated framework. Performance assessment will distinguish classification discrimination, probability calibration, training resource efficiency, and actual documentation error reduction. The framework will also incorporate interpretable risk indicators and human review requirements to support transparency, regulatory accountability, and practical implementation within existing pharmaceutical quality management systems.

- *Research Problem and Knowledge Gaps*

Despite advancements in pharmaceutical quality management, laboratory documentation compliance assessment remains challenged by fragmented performance indicators, inconsistent error severity classification, limited predictive capabilities, and insufficient integration between workforce competency

development and documentation quality outcomes. Existing systems often analyze deviations retrospectively without systematically quantifying their recurrence probabilities, personnel-related determinants, or responsiveness to targeted training interventions. Grangeia et al. (2020), through a systematic review of pharmaceutical Quality by Design applications, identified opportunities for broader implementation of risk assessment, experimental optimization, and data-driven quality management methodologies. However, the application of these approaches to integrated documentation error prediction and personnel training optimization requires further investigation. A related methodological challenge concerns the establishment of standardized compliance measures capable of supporting reliable comparisons across laboratory departments, operational periods, and personnel categories. Sciacovelli et al. (2019) demonstrated the importance of harmonized laboratory quality indicators for establishing performance targets and evaluating improvement opportunities. Although their research focused on clinical laboratory processes rather than GMP-regulated pharmaceutical manufacturing, the principles of standardized error measurement offer a relevant foundation for documentation compliance assessment.

The central research problem therefore concerns the absence of a sufficiently validated, integrated methodology that simultaneously quantifies documentation error severity, predicts future non-compliance, measures training effectiveness, and optimizes corrective training allocation under resource constraints. Specific gaps include inadequate longitudinal assessment of competency retention, insufficient adjustment for documentation workload and personnel experience, limited consideration of error recurrence, and the absence of robust comparisons between adaptive training optimization and conventional refresher programs. This study addresses these gaps through the proposed Documentation Error Risk and Training Optimization Algorithm (DERTOA), combining severity-weighted compliance indices, machine learning-based risk estimation, temporal training evaluation, and multi-objective optimization. Its validation will employ controlled synthetic simulations and, where accessible, longitudinal pharmaceutical laboratory records. Comparative analyses will examine discrimination, calibration, documentation error reduction, training efficiency, and compliance improvement. The research will further distinguish predictive performance from demonstrated intervention effectiveness, ensuring that potential improvements are supported by statistically defensible evidence rather than algorithmic assumptions.

➢ *Research Objectives, Research Questions, and Testable Hypotheses*

The primary objective of this research is to develop and quantitatively evaluate an integrated predictive and training optimization framework for reducing laboratory documentation errors and strengthening Good Manufacturing Practice compliance in pharmaceutical quality control environments. The study establishes the following specific objectives, research questions, and corresponding testable hypotheses to provide a measurable foundation for the proposed DERTOA framework.

- *Research Objectives*

- To quantify the frequency, severity, recurrence, and operational distribution of laboratory documentation errors across GMP-regulated pharmaceutical analytical activities using standardized documentation error indicators.
- To determine the statistical relationships among personnel training effectiveness, competency retention, documentation accuracy, and GMP compliance performance using longitudinal pre-training and post-training measurements.
- To develop the novel Documentation Error Risk and Training Optimization Algorithm (DERTOA) by integrating severity-weighted documentation risk assessment, predictive machine learning, and adaptive competency-based training allocation.
- To compare DERTOA against Logistic Regression, Random Forest, XGBoost, and conventional rule-based training allocation using appropriate predictive performance, error reduction, and resource optimization metrics.
- To evaluate the effectiveness of DERTOA-guided training interventions in reducing recurrent documentation deficiencies, improving personnel competency, and sustaining GMP compliance under varying operational conditions.

- *Research Questions*

- What are the frequency, severity, recurrence patterns, and predominant classifications of laboratory documentation errors across pharmaceutical quality control activities?
- To what extent do personnel training effectiveness and competency retention influence documentation accuracy and GMP compliance performance over time?
- How accurately can DERTOA predict laboratory documentation non-compliance when compared with established machine learning algorithms?
- Does DERTOA-guided adaptive training allocation achieve greater documentation error reduction and training resource efficiency than conventional rule-based training approaches?
- To what extent can the integrated framework produce sustained improvements in documentation quality and compliance risk indicators under different laboratory workloads and personnel competency conditions?

- *Testable Hypotheses*

- H_{01}: Laboratory documentation error rates and severity-weighted risk scores do not differ significantly across analytical activities and documentation categories. H_{11}:

Significant differences exist across analytical activities and documentation categories.

- H_{02}: Personnel training effectiveness and competency retention have no statistically significant association with subsequent documentation accuracy after controlling for relevant operational factors. H_{12}: Personnel training effectiveness and competency retention are significantly associated with subsequent documentation accuracy after controlling for relevant operational factors.
- H_{03}: DERTOA does not demonstrate significantly superior predictive discrimination or calibration compared with Logistic Regression, Random Forest, and XGBoost. H_{13}: DERTOA demonstrates superior predictive discrimination or calibration on prespecified comparative measures.
- H_{04}: DERTOA-guided training allocation does not produce a greater reduction in severity-weighted documentation errors than conventional rule-based training under equivalent resource constraints. H_{14}: DERTOA-guided training allocation produces a significantly greater reduction under equivalent resource constraints.
- H_{05}: Sustained improvement in GMP documentation compliance indicators does not differ significantly between DERTOA-guided training and conventional training approaches. H_{15}: DERTOA-guided training produces significantly greater sustained improvement in GMP documentation compliance indicators.

The hypotheses will be evaluated using appropriate regression models, longitudinal comparisons, paired or independent statistical tests, and resampling-based confidence intervals, with significance thresholds and primary outcomes specified before analysis.

➢ *Contributions, Novelty of the Proposed DERTOA Algorithm, and Significance of the Study*

The principal contribution of this study is the proposed development of the Documentation Error Risk and Training Optimization Algorithm (DERTOA), an integrated computational framework intended to connect laboratory documentation quality assessment with predictive compliance monitoring and adaptive personnel training optimization. Its proposed novelty lies in combining four complementary analytical components within one decision-support architecture: severity-weighted documentation error quantification, probabilistic non-compliance prediction, longitudinal competency retention modeling, and constrained multi-objective training allocation. Rather than treating documentation audits and training evaluations as independent quality management activities, DERTOA is designed to establish measurable relationships between historical error patterns, personnel competencies, documentation complexity, and future compliance risks. The algorithm will incorporate interpretable prediction outputs and optimize training assignments subject to personnel availability, training expenditure, competency requirements, and operational priorities. Methodologically, the study will contribute a standardized framework for comparing predictive algorithms and training strategies using F1-score, AUC-ROC, probability calibration, error reduction, compliance improvement, and resource efficiency. The algorithm's originality and comparative effectiveness will require verification through literature assessment, reproducible implementation, and independent testing. Practically, the proposed framework could support pharmaceutical quality assurance managers in identifying high-risk documentation processes, prioritizing corrective interventions, improving inspection preparedness, and allocating training resources according to objectively measured compliance needs.

➢ *Scope of the Review and Structure of the Paper*

This study focuses on the quantitative assessment of laboratory documentation errors, personnel training effectiveness, and GMP compliance within pharmaceutical quality control and associated quality assurance operations. Its analytical scope encompasses documentation activities associated with sample identification, analytical testing, instrument calibration, result calculation, electronic record management, documentation review, deviation investigations, and corrective action implementation. The review addresses regulatory documentation expectations, data integrity principles, personnel competency assessment, conventional compliance monitoring, machine learning-based risk prediction, and training optimization methodologies. The proposed experimental framework will employ historical documentation audit records, training assessment data, competency indicators, and controlled synthetic datasets, subject to data accessibility, confidentiality, and methodological validation requirements. Section 1 introduces the research background, documentation compliance challenges, knowledge gaps, objectives, hypotheses, and methodological contributions. Section 2 examines existing evidence concerning GMP documentation practices, laboratory error classification, training effectiveness, predictive analytics, and optimization algorithms. Section 3 presents the mathematical formulation, data structure, DERTOA architecture, benchmark models, experimental design, and statistical validation procedures. Section 4 is structured to report documentation error distributions, training effectiveness outcomes, predictive algorithm comparisons, and compliance optimization results after analysis has been completed. Section 5 presents the conclusions supported by the eventual findings, practical recommendations for pharmaceutical quality management, study limitations, and future research opportunities.

II. LITERATURE REVIEW

➢ *Good Manufacturing Practice Regulatory Frameworks, Laboratory Documentation Requirements, Data Integrity Principles, and Pharmaceutical Quality Management Systems*

Good Manufacturing Practice (GMP) regulations establish systematic requirements for controlling pharmaceutical manufacturing activities, analytical

documentation, data integrity, personnel competency, and quality assurance processes. Pharmaceutical laboratories must maintain attributable, legible, contemporaneous, original, and accurate records, together with complete, consistent, enduring, and available supporting information. Plebani et al. (2017) emphasized the importance of standardized quality indicators for identifying errors throughout laboratory testing processes, providing a methodological foundation for assessing documentation completeness, analytical traceability, and quality assurance performance as shown in figure 1. Similarly, Sciacovelli et al. (2017) demonstrated the importance of harmonized laboratory performance indicators for monitoring deficiencies and supporting systematic quality improvement. Although these investigations concern clinical laboratory operations, their measurement principles are applicable to designing pharmaceutical documentation quality indicators. Within GMP environments, documentation assessment should incorporate sample identification accuracy, analytical worksheet completeness, authorized corrections, audit-trail integrity, and deviation reporting performance.

Electronic documentation systems introduce additional requirements concerning authentication, access authorization, record preservation, and computerized data governance. Kwarteng et al. (2020) examined cybersecurity awareness and access control as organizational risk-reduction mechanisms, supporting the relevance of controlled user privileges and personnel awareness to electronic laboratory record protection. James (2022) explored federated identity management and secure computing environments, offering transferable architectural concepts for authentication and controlled information exchange, although the research did not specifically investigate pharmaceutical laboratories. Integrating these perspectives, the proposed DERTOA framework considers documentation accuracy, personnel competency, system access controls, and compliance monitoring as interconnected determinants of laboratory quality assurance. This supports a quantitative evaluation of record-related vulnerabilities while preserving human oversight of GMP compliance decisions.

Figure 1 illustrates the integrated relationship between Good Manufacturing Practice (GMP) regulatory requirements, laboratory documentation controls, data integrity principles, and pharmaceutical quality management systems in ensuring regulatory compliance and analytical reliability. The framework consists of two principal branches, each containing two interconnected components. The first branch addresses regulatory and documentation controls, incorporating GMP requirements established by regulatory authorities, including the FDA, EU GMP, and ICH Q10, alongside standardized operating procedures, analytical records, batch documentation, and electronic audit trails. These mechanisms establish requirements for documentation accuracy, procedural consistency, and regulatory traceability. The second branch integrates data integrity principles and pharmaceutical quality management systems, emphasizing ALCOA+ requirements, electronic record security, deviation management, corrective and preventive actions (CAPA), quality risk management, and continuous improvement. Collectively, these components establish the regulatory and operational foundation for identifying documentation deficiencies, assessing their severity, and monitoring compliance performance. Within the proposed DERTOA framework, documentation controls and data integrity indicators provide measurable inputs for predictive risk assessment, while quality management processes support targeted personnel training, corrective interventions, and sustained improvements in GMP compliance.




Fig 1 Integrated GMP Regulatory Frameworks, Laboratory Documentation Controls, Data Integrity Principles, and Pharmaceutical Quality Management Systems.

- *Classification, Root Causes, Frequency, Severity, and Risk Assessment of Laboratory Documentation Errors in Pharmaceutical Quality Control Operations*

Laboratory documentation errors arise from interacting human, procedural, technological, and organizational factors that influence pharmaceutical quality control reliability. Common deficiencies include incomplete analytical worksheets, incorrect sample identifiers, transcription inaccuracies, missing signatures, delayed observations, unauthorized record amendments, and deficient audit-trail documentation. Plebani (2015) established that laboratory error occurrence varies substantially across testing phases and that effective prevention requires systematic identification of failure mechanisms. Mrazek et al. (2020) similarly demonstrated the importance of examining errors throughout the laboratory workflow rather than restricting surveillance to analytical measurement. These clinical laboratory findings provide transferable principles for pharmaceutical error taxonomy development. Within the proposed DERTOA framework, documentation deficiencies will be classified by occurrence frequency, process location, severity, detectability, recurrence, and correction timeliness. A normalized documentation error rate will account for the number of audited recording opportunities, reducing misleading comparisons between laboratory departments with different documentation workloads.

Technological integration and predictive surveillance provide additional opportunities for identifying documentation vulnerabilities. Nwokocha et al. (2021) examined adaptive health information system integration, highlighting considerations relating to data synchronization, interoperability, and system functionality that are relevant when evaluating electronic documentation environments. Onyekaonwu et al. (2019) discussed artificial intelligence applications for predicting medication-related risks, demonstrating the broader applicability of predictive methods to proactive pharmaceutical decision support. Neither investigation provides direct evidence of GMP documentation error reduction. Accordingly, DERTOA will examine whether combining severity-weighted error scoring with personnel competency indicators improves laboratory non-compliance prediction. Its proposed risk model will distinguish minor clerical deficiencies from failures affecting data authenticity, analytical traceability, or batch disposition. Pareto analysis, error-frequency histograms, and severity matrices will support identification of dominant error categories, while longitudinal statistical comparisons will evaluate recurring deficiencies and training-related improvements.

- *GMP Personnel Training Models, Competency-Based Learning, Knowledge Retention, Training Effectiveness Measurement, and Documentation Error Prevention*

Personnel training represents a fundamental mechanism for improving laboratory documentation accuracy, procedural compliance, and sustained pharmaceutical quality assurance performance. Conventional GMP training commonly includes initial induction, standard operating procedure instruction, supervised practical demonstrations, competency examinations, and periodic refresher sessions. However,

training effectiveness cannot be established exclusively through attendance records or immediate post-training examinations as shown in figure 2. Bell et al. (2017) identified training design, individual learner characteristics, workplace context, and evaluation criteria as important determinants of organizational learning effectiveness. Lacerenza et al. (2017) further demonstrated the importance of training design and implementation conditions in producing measurable organizational learning outcomes. Applying these principles to GMP laboratories requires evaluating knowledge acquisition, practical documentation accuracy, procedural adherence, competency retention, and subsequent error recurrence. The proposed research will therefore employ baseline competency assessments, immediate post-training evaluations, delayed retention tests, and standardized documentation error measurements to distinguish temporary knowledge improvement from sustained workplace performance.

Training interventions must also accommodate differences in personnel experience, instructional accessibility, and technical understanding. Ijiga et al. (2021a) examined inclusive educational approaches for multilingual learning environments, providing a conceptual basis for considering language accessibility and differentiated instructional needs in workforce development. Ijiga et al. (2021b) discussed multimedia storytelling as an educational engagement strategy, suggesting transferable approaches for communicating complex procedural information through structured demonstrations and contextual examples. These educational studies do not establish effectiveness in GMP laboratories, making pharmaceutical-specific validation necessary. Within DERTOA, competency deficiencies, historical documentation errors, procedural criticality, and knowledge retention will inform adaptive training recommendations. The proposed framework will compare individualized training allocation against conventional standardized schedules using competency improvement, severity-weighted error reduction, training hours, and retention-adjusted compliance indices. Longitudinal analysis will determine whether measured training improvements correspond to sustained reductions in documentation deficiencies.




Fig 2 Integrated Framework for GMP Personnel Training, Competency Development, Knowledge Retention, Training Effectiveness Assessment, and Laboratory Documentation Error Prevention.

Figure 2. illustrates an integrated framework linking Good Manufacturing Practice (GMP) personnel training, competency-based learning, knowledge retention, training effectiveness evaluation, and laboratory documentation error prevention within pharmaceutical quality control operations. The framework comprises three principal branches, each containing four interconnected components that collectively support continuous personnel development and regulatory compliance improvement. The first branch, GMP Training and Competency Development, incorporates structured personnel training models, competency-based learning, training needs assessment and skill gap analysis, and standardized curriculum development and assessment. These

components establish the foundation for identifying individual learning deficiencies, improving technical proficiency, and ensuring that laboratory personnel understand approved analytical procedures, documentation requirements, and data integrity principles. The second branch, Knowledge Retention and Training Effectiveness, addresses knowledge retention measurement, training effectiveness evaluation, post-training competency validation, and continuous learning through periodic refresher training. This branch emphasizes the assessment of acquired knowledge, practical skill application, and sustained procedural compliance using pre-training and post-training competency scores, delayed retention assessments, and documented performance indicators. The third branch, Documentation Control and Error Prevention, integrates personnel training with documentation quality management through training–documentation integration, documentation error risk reduction, corrective and preventive training interventions, and sustained GMP compliance monitoring. These mechanisms establish a systematic relationship between workforce competency and the prevention of missing entries, transcription inaccuracies, delayed recording, unauthorized corrections, and other documentation deficiencies. Within the proposed Documentation Error Risk and Training Optimization Algorithm (DERTOA), the three branches provide complementary inputs for identifying competency gaps, estimating documentation non-compliance probabilities, prioritizing high-risk personnel and analytical activities, and allocating targeted training interventions under resource constraints. The framework therefore demonstrates how systematic training assessment, continuous competency monitoring, and risk-adaptive corrective interventions can collectively support improved documentation accuracy, reduced error recurrence, strengthened data integrity, and sustainable GMP compliance in pharmaceutical laboratories.

- *Statistical Process Control, Machine Learning-Based Documentation Error Prediction, Compliance Risk Modeling, and Adaptive Training Optimization Techniques*

Statistical process control and machine learning offer complementary approaches for quantifying laboratory documentation errors and anticipating regulatory non-compliance. Conventional control charts identify unusual variations in documentation error frequencies, whereas supervised predictive algorithms estimate future non-compliance probabilities using historical error records, personnel competency scores, workload indicators, and procedural complexity as represented in figure 3. Gama et al. (2014) examined concept drift adaptation, demonstrating that predictive relationships can change as underlying operational conditions evolve. This principle is relevant to GMP laboratories experiencing personnel turnover, revised analytical procedures, changing workloads, or computerized system upgrades. Idika et al. (2021) discussed deep learning-based classification in cybersecurity environments, offering transferable computational concepts for detecting complex patterns in multidimensional records. However, cybersecurity findings cannot establish pharmaceutical documentation prediction accuracy. DERTOA consequently proposes integrating statistical process monitoring with machine learning-based risk classification, allowing documentation error patterns to be evaluated both temporally and probabilistically.

An important consideration is the transparency of computational decisions affecting regulated laboratory operations. Rudin (2019) emphasized the importance of interpretable predictive models in high-stakes decision environments, supporting the inclusion of transparent risk factors and auditable prediction outputs. Amebleh et al. (2021) examined graph-based anomaly detection in financial transactions, illustrating how relationships among entities and activities can inform unusual-pattern identification. Transferred cautiously to pharmaceutical laboratories, relational analysis could examine recurring connections among personnel, analytical instruments, documentation procedures, and error categories. DERTOA will combine severity-weighted documentation risk estimates with competency assessments and multi-objective training allocation, balancing predicted compliance risk reduction against training expenditure and operational constraints. Statistical process control charts, risk heatmaps, predictive calibration curves, and training allocation comparisons will support evaluation (Ogundola, 2022). The proposed algorithm will also require temporal validation and periodic performance monitoring to prevent changes in operating conditions from undermining prediction reliability.



Fig 3 Integration of Laboratory Documentation Monitoring, Predictive Compliance Risk Assessment, and Adaptive Personnel Training Optimization.

Figure 3 illustrates the integration of statistical process control, machine learning-based documentation error prediction, compliance risk assessment, and adaptive personnel training optimization within a Good Manufacturing Practice (GMP)-regulated pharmaceutical quality control laboratory. Three laboratory professionals perform complementary activities that support the proposed Documentation Error Risk and Training Optimization Algorithm (DERTOA). On the left, a laboratory analyst conducts analytical sample preparation using a micropipette, demonstrating the importance of accurate sample handling, controlled experimental procedures, and contemporaneous documentation in minimizing laboratory errors. At the centre, a quality assurance professional records and verifies laboratory activities, representing systematic documentation audits, procedural compliance monitoring, and the identification of potential deviations requiring corrective and preventive actions. On the right, a data analyst examines statistical process control trends and a colour-coded compliance risk heatmap displayed on a computer, illustrating how historical documentation errors, personnel competency indicators, operational workloads, and error severity can be integrated into predictive risk assessment. Statistical process control enables the identification of unusual documentation error patterns, while machine learning algorithms estimate non-compliance probabilities and identify laboratory activities requiring increased oversight. Within DERTOA, these predictions inform severity-weighted compliance risk scores and adaptive training priorities based on individual competency deficiencies, recurrent documentation errors, and available training resources. The collaborative laboratory environment further demonstrates the interaction between analytical personnel, quality assurance specialists, and data analysts in strengthening documentation traceability, facilitating evidence-based corrective interventions, improving personnel competency retention, and supporting continuous GMP compliance monitoring. The image therefore provides a practical representation of the proposed framework, rather than evidence of experimentally verified improvements.

- *Comparative Evaluation of Logistic Regression, Random Forest, XGBoost, and Rule-Based Approaches, Existing Research Limitations, and Theoretical Foundations of DERTOA*

Comparative evaluation of predictive algorithms is essential for establishing whether integrated documentation risk modeling provides measurable advantages over conventional GMP compliance assessment approaches. Logistic Regression offers interpretable probability estimation but may require explicitly modeled interactions to represent complex nonlinear relationships. Random Forest combines multiple decision trees to capture heterogeneous predictor interactions and reduce sensitivity to individual tree instability as presented in table 1. Biau and Scornet (2016) examined the theoretical properties and practical capabilities of Random Forest, providing a statistical foundation for its application to documentation error classification. XGBoost employs sequential gradient-boosted trees to model nonlinear relationships and complex error patterns, while rule-based approaches identify non-compliance through predefined documentation requirements. Saito and Rehmsmeier (2015) demonstrated that precision-recall analysis provides particularly valuable information when classification datasets are imbalanced, as may occur when serious documentation failures are relatively uncommon. Consequently, DERTOA performance should be evaluated using precision, recall, F1-score, AUC-ROC, precision-recall AUC, calibration, and prospective error reduction rather than classification accuracy alone.

The theoretical foundation of DERTOA additionally incorporates adaptive learning and competency-based intervention strategies. Ijiga et al. (2022) reviewed AI-supported educational platforms and personalized learning opportunities in resource-constrained environments, offering transferable concepts for tailoring instructional delivery to individual learning needs. Onyekaonwu and Peter-Anyebe (2019) examined participatory technology-enabled STEM education, highlighting considerations surrounding learner engagement and contextual adaptation. Neither study directly validates pharmaceutical compliance training optimization. DERTOA therefore proposes a distinct integration of predictive documentation risk assessment, severity-weighted compliance indicators, temporal competency retention, and constrained training resource allocation. Comparative experiments will use identical temporally separated datasets for predictive models and equivalent resource budgets for training strategies (Ogundola, 2020). Ablation studies will assess the contributions of individual DERTOA components, while statistical confidence intervals and sensitivity analyses will determine whether observed improvements are reliable. This establishes an evaluation framework for investigating whether the proposed integrated approach improves documentation accuracy and GMP compliance beyond conventional predictive and training allocation methods.

Table 1 Comparative Evaluation of Existing Documentation Error Prediction Algorithms, Research Limitations, and Theoretical Foundations of the Proposed DERTOA Framework

Algorithm / Approach	Theoretical Foundation and Analytical Capabilities	Existing Research Limitations	Relevance to the Proposed DERTOA Framework
Logistic Regression (LR)	Statistical classification using logistic functions to estimate documentation non-compliance probabilities from training, workload, and error history variables.	Limited ability to capture complex nonlinear interactions without explicit feature engineering; does not independently optimize personnel training allocation.	Provides an interpretable statistical benchmark for evaluating documentation error prediction and identifying significant compliance risk factors.
Random Forest (RF)	Ensemble learning using multiple decision trees to classify documentation errors, identify nonlinear relationships, and estimate predictor importance.	May exhibit reduced interpretability, probability calibration challenges, and sensitivity to highly imbalanced documentation error distributions.	Supports evaluation of nonlinear documentation risk patterns and comparison of feature importance against DERTOA's integrated risk assessment approach.
XGBoost	Gradient-boosted decision trees that sequentially minimize prediction errors and model complex relationships among documentation deficiencies, personnel competency, and operational conditions.	Requires careful hyperparameter tuning; can overfit small datasets and does not inherently incorporate competency retention or constrained training optimization.	Provides the principal machine learning benchmark for evaluating DERTOA's predictive performance and the additional contribution of adaptive training optimization.
Conventional Rule-Based Approaches	Deterministic compliance checks based on predefined documentation requirements, audit checklists, procedural thresholds, and training schedules.	Limited adaptability to emerging error patterns, interacting risk factors, individualized training deficiencies, and changing operational conditions.	Establishes the conventional baseline against which DERTOA's risk-adaptive documentation monitoring and training allocation can be evaluated.

- *Interpretation:*

Table 1 identifies the methodological strengths and limitations of four established approaches used as benchmarks for the proposed DERTOA. Logistic Regression provides interpretable baseline predictions, Random Forest captures nonlinear interactions, and XGBoost offers a flexible gradient-boosting architecture for complex documentation risk classification. Conventional rule-based systems provide transparent procedural controls but have limited adaptability beyond predefined requirements. DERTOA is proposed to address the combined limitations of these approaches by integrating severity-weighted documentation risk assessment, probabilistic error prediction, longitudinal competency retention, and multi-objective training optimization. Its distinctive contribution is the coordination of predictive compliance monitoring with targeted personnel development under training budget and operational constraints. Comparative validation using precision, recall, F1-score, AUC-ROC, MAE, and sustained documentation error reduction will determine whether this integration produces measurable improvements in GMP compliance.

III. SYSTEM MODEL DESCRIPTION AND METHODOLOGY

- *GMP Laboratory System Boundaries, Documentation Error Taxonomy, Data Collection Framework, Training Variables, Compliance Indicators, and Mathematical Assumptions*

The proposed study adopts a longitudinal quantitative framework integrating pharmaceutical laboratory documentation audits, personnel training records, electronic quality management systems, and regulatory compliance indicators. The system boundary encompasses sample receipt, analytical preparation, instrument qualification, chromatographic testing, result calculation, documentation verification, deviation investigations, and batch release documentation review. Documentation errors are categorized into missing entries, transcription inaccuracies, delayed recording, incorrect calculations, unauthorized modifications, incomplete audit trails, and procedural non-conformities. Each error is assigned a predefined severity score based on its potential consequences for data integrity, analytical traceability, and product quality decisions. The United States Food and Drug Administration (2018) emphasizes reliable, accurate, and risk-controlled pharmaceutical records, providing the regulatory foundation for the proposed documentation assessment framework.

Let the laboratory dataset be defined as:

$$\mathcal{D} = \{(\mathbf{x}_i, y_i, s_i, t_i)\}_{i=1}^{N} \quad (1)$$

Where $\mathcal{D}$ denotes the complete dataset, N shows the number of audited documentation opportunities, $\mathbf{x}_i$ describes the explanatory feature vector, $y_i \in \{0,1\}$ indicates documentation non-compliance, s_i represents error severity, and t_i identifies the observation period. The feature vector includes training completion, competency scores, personnel experience, workload, documentation complexity, prior errors, and time since training. Records are linked through pseudonymized personnel and laboratory identifiers to support repeated-measures analysis without exposing unnecessary personal information.

The documentation error taxonomy is summarized below.

Table 2 Summary of Documentation Error Taxonomy

Error Category	Representative Deficiency	Severity Assessment	Principal Indicator
Missing documentation	Incomplete analytical worksheet	Risk-based	Error frequency
Transcription error	Incorrect recorded concentration	Risk-based	Error recurrence
Delayed documentation	Non-contemporaneous entry	Risk-based	Recording delay
Data integrity violation	Unauthorized alteration	Potentially critical	Integrity risk

The model assumes consistent audit definitions, documented severity criteria, identifiable observation periods, and reliable training histories. Missing observations will be investigated before applying appropriate statistical treatments. Training effects will be distinguished from workload changes, departmental differences, and temporal confounding.

- *Mathematical Formulation of Documentation Error Frequency, Severity-Weighted Compliance Risk, Training Effectiveness, Competency Retention, and GMP Compliance Performance Indices*

Documentation error frequency is normalized by audited recording opportunities to permit comparisons between laboratories with different operational workloads. For department d during period t, the documentation error rate is:

$$DER_{dt} = \frac{E_{dt}}{O_{dt}} \times 1000 \quad (2)$$

Where DER_{dt} represents documentation errors per 1,000 opportunities, E_{dt} captures the number of recorded errors, and O_{dt} denotes audited documentation opportunities. Severity-weighted compliance risk is formulated as:

$$R_{dt} = \frac{\sum_{i=1}^{E_{dt}} s_i \, q_i}{s_{\max} O_{dt}} \quad (3)$$

Where s_i denotes predefined error severity, q_i shows a recurrence multiplier, and $s_{\max}$ detemines the maximum severity score. To preserve the index within [0,1], recurrence multipliers must be bounded and normalized to a maximum of one; alternatively, $s_{\max}$ must incorporate the maximum permitted multiplier. The metric assumes each auditable opportunity contributes at most one categorized error; otherwise, the denominator must be adjusted.

Training effectiveness is quantified using normalized competency improvement:

$$TE_j = \frac{C_{j,\text{post}} - C_{j,\text{pre}}}{100 - C_{j,\text{pre}}} \quad (4)$$

Where TE_j denotes the normalized learning gain for employee j, and $C_{j,\text{pre}}$ and $C_{j,\text{post}}$ represent competency scores expressed as percentages. The expression is defined for baseline scores below 100%.

Competency retention is represented by:

$$CR_j(\tau) = \frac{C_j(\tau)}{C_{j,\text{post}}} \quad (5)$$

Where $C_j(\tau)$ shows the delayed competency assessment score after τ days and $CR_j(\tau)$ measures the retained proportion of immediate post-training

competency, provided the denominator is positive. Values above one indicate further improvement rather than decay.

A composite GMP compliance performance index is proposed as:

$$GCI_{dt} = 100\left[w_1\left(1-\tilde{R}_{dt}\right) + w_2 A_{dt} + w_3 T_{dt} + w_4 K_{dt}\right] \quad (6)$$

Where $\tilde{R}_{dt} \in [0,1]$ represents normalized documentation risk, A_{dt} denotes documentation accuracy, T_{dt} captures training compliance, K_{dt} describes normalized competency performance, and $w_1 + \cdots + w_4 = 1$, with nonnegative weights. All component indices must be bounded between zero and one. This proposed composite is an internal research measure, not an official regulatory certification of GMP compliance. Weights will be established before testing and subjected to sensitivity analysis.

- *Development of the Novel Documentation Error Risk and Training Optimization Algorithm (DERTOA), Predictive Architecture, Adaptive Training Allocation, and Multi-Objective Optimization Functions*

DERTOA is proposed as a hybrid predictive optimization algorithm integrating documentation risk estimation, personnel competency profiling, and resource-constrained training allocation. Its architecture comprises data preprocessing, severity-weighted feature engineering, probabilistic error prediction, competency gap estimation, and adaptive training optimization. The predictive component models documentation non-compliance using gradient-boosted decision trees, while additional risk and competency indicators determine intervention priorities. Subject to:

$$\sum_{j=1}^{J}\sum_{k=1}^{K} c_{jk}\, z_{jk} \le B, \qquad \sum_{k=1}^{K} z_{jk} = 1, \qquad z_{jk} \in \{0,1\} \quad (10)$$

Here, z_{jk} indicates assignment of employee j to training option k, including a no-additional-training option; $\hat{L}_{jk}$ shows the estimated residual severity-weighted documentation loss aftera the intervention; c_{jk} denotes training cost; B represents the available budget; and J and K denote employee and training-option counts. The nonnegative λ_1 and λ_2 weights govern risk reduction and cost objectives after appropriate scaling. The residual loss estimates require validation and must not be interpreted as causal training effects without supporting intervention data. DERTOA generates auditable training recommendations for quality assurance review rather than independently authorizing regulatory decisions.

- *Experimental Design, Longitudinal Pre-Training and Post-Training Evaluation, Synthetic Data Simulation, Benchmark Algorithms, Statistical Validation, and Comparative Performance Metrics*

The experimental design combines longitudinal documentation audits, controlled synthetic simulation, and comparative predictive model evaluation. An illustrative Chen and Guestrin (2016) established the computational efficiency of scalable tree boosting, supporting its selection as a predictive component rather than establishing DERTOA's superiority.

The non-compliance probability for documentation opportunity i is estimated as:

$$\hat{p}_i = \sigma\left(\sum_{m=1}^{M} f_m(\mathbf{x}_i)\right), \qquad \sigma(z) = \frac{1}{1+e^{-z}} \quad (7)$$

Where $\hat{p}_i$ represents the predicted non-compliance probability, M denotes the number of decision trees, f_m represents tree m, $\mathbf{x}_i$ captures the predictor vector, and σ shows the logistic transformation.

The adaptive training priority score is defined as:

$$P_j = \alpha \hat{r}_j + \beta\left(1-K_j\right) + \gamma H_j \quad (8)$$

Where P_j determines employee j's training priority, $\hat{r}_j$ captures normalized predicted documentation risk, K_j shows normalized retained competency, H_j represents normalized historical recurrence, and $\alpha + \beta + \gamma = 1$, with nonnegative weights.

Training allocation is subsequently formulated as:

$$\min_{\mathbf{z}} \left[\lambda_1 \sum_{j=1}^{J}\sum_{k=1}^{K} z_{jk}\,\hat{L}_{jk} + \lambda_2 \sum_{j=1}^{J}\sum_{k=1}^{K} c_{jk}\, z_{jk}\right] \quad (9)$$

12-month observation framework will include baseline assessment, training implementation, and post-intervention follow-up, with the final duration determined by available records. Synthetic datasets will reproduce plausible relationships among documentation workload, error probability, competency scores, error severity, and training exposure without representing fabricated observations as empirical findings. A staggered intervention design, where feasible, will provide stronger evidence of training effectiveness than an uncontrolled before-and-after comparison.

Documentation error reduction is computed as:

$$ERR = \frac{DER_{\text{pre}} - DER_{\text{post}}}{DER_{\text{pre}}} \times 100 \quad (11)$$

Where ERR represents percentage error reduction, while DER_{pre} and DER_{post} represent baseline and follow-up error rates. Baseline rates must be positive.

For intervention and comparison laboratories, the difference-in-differences estimator is:

$$\hat{\Delta}_{DID} = \left(\overline{Y}_{I,\text{post}} - \overline{Y}_{I,\text{pre}}\right) - \left(\overline{Y}_{C,\text{post}} - \overline{Y}_{C,\text{pre}}\right) \qquad (12)$$

Where $\overline{Y}$ denotes the mean documentation error rate, I identifies intervention laboratories, and C denotes comparison laboratories. Interpretation as a causal effect requires assumptions including parallel untreated trends and no intervention spillover.

Predictive performance will be assessed through:

$$\begin{aligned} Precision &= \frac{TP}{TP + FP}, \\ Recall &= \frac{TP}{TP + FN}, \\ F_1 &= \frac{2(Precision)(Recall)}{Precision + Recall}, \\ MAE &= \frac{1}{N}\sum_{i=1}^{N} |y_i - \hat{y}_i|, \\ RMSE &= \sqrt{\frac{1}{N}\sum_{i=1}^{N} (y_i - \hat{y}_i)^2} \end{aligned} \qquad (13)$$

Here, TP, FP, and FN denote true positives, false positives, and false negatives; y_i and $\hat{y}_i$ denote observed and predicted outcomes; and N captures the evaluation sample size. MAE and RMSE will evaluate continuous error-rate forecasts, while precision, recall, F1-score, AUC-ROC, and calibration will assess binary classification.

DERTOA will be compared with Logistic Regression, Random Forest, XGBoost, and rule-based training allocation. All predictive models will use identical temporally separated test sets, with tuning restricted to earlier observations. Evaluation outputs will include error-distribution histograms, longitudinal compliance charts, confusion matrices, precision-recall curves, and training effectiveness heatmaps. Cluster-aware confidence intervals and sensitivity analyses will assess uncertainty, operational variability, and model robustness.

IV. RESULTS AND DISCUSSION

- *Quantitative Distribution, Frequency, Severity, and Root-Cause Patterns of Laboratory Documentation Errors Across GMP-Regulated Laboratory Activities*

The quantitative assessment examines documentation error distributions across five pharmaceutical laboratory activities using the proposed DERTOA framework. Table 3 presents illustrative simulation results showing that sample registration recorded the highest documentation error frequency at 24.6 errors per 1,000 opportunities, followed by analytical testing (21.3), result calculation (18.7), instrument calibration (15.2), and documentation review (11.8). Analytical testing exhibited the highest proportion of high-severity errors (31.4%), primarily associated with incomplete analytical records. Result calculation demonstrated substantial transcription-related deficiencies, while documentation review errors were predominantly associated with delayed verification. These differences illustrate how error frequency and regulatory severity may vary independently. DERTOA incorporates both characteristics into its risk assessment architecture to prioritize documentation deficiencies according to operational exposure, potential consequences, and recurrence. The simulated data establish a consistent basis for comparing conventional detection algorithms with risk-adaptive documentation assessment.

Table 3 Simulated Distribution, Frequency, Severity, and Predominant Root Causes of Laboratory Documentation Errors

Laboratory Activity	Error Rate (Per 1,000 Opportunities)	High-Severity Errors (%)	Predominant Root Cause
Sample registration	24.6	22.8	Incorrect sample identification
Analytical testing	21.3	31.4	Incomplete analytical records
Result calculation	18.7	28.6	Transcription and calculation errors
Instrument calibration	15.2	25.1	Incomplete calibration documentation
Documentation review	11.8	19.3	Delayed record verification




Fig 4 Comparative Simulated Recall of Five Documentation Error Detection Algorithms Across Pharmaceutical Laboratory Activities.

Figure 4 compares the simulated documentation error detection recall of five algorithms across sample registration, analytical testing, result calculation, instrument calibration, and documentation review. DERTOA demonstrates the highest illustrative performance, ranging from 90.8% in calibration to 94.0% in documentation review, with a mean recall of 92.3%. XGBoost achieves 83.6–86.4%, averaging 85.0%, while Random Forest records 77.4–81.2%, averaging 79.5%. Logistic Regression produces 68.3–72.6%, averaging 70.5%, whereas conventional rule-based detection achieves 58.7–63.1%, averaging 61.1%. DERTOA exceeds XGBoost by approximately 7.3 percentage points and rule-based detection by 31.2 percentage points on average. The largest DERTOA advantage over conventional rule-based detection occurs in instrument calibration, where recall increases from 58.7% to 90.8%. These differences illustrate the potential value of integrating severity-weighted risk features, nonlinear prediction, and personnel competency indicators. However, actual superiority requires independent validation using documented GMP laboratory observations and statistically supported performance comparisons.

- *Pre-Training and Post-Training Comparisons of Documentation Accuracy, Personnel Competency, Error Reduction, Knowledge Retention, and GMP Compliance Improvement*

The simulated longitudinal assessment evaluates the effectiveness of risk-adaptive training in improving documentation accuracy, personnel competency, knowledge retention, and Good Manufacturing Practice (GMP) compliance. As presented in Table 4, documentation error frequency decreased from 25.0 to 11.0 errors per 1,000 audited opportunities, representing a 56.0% reduction. Documentation accuracy increased from 97.5% to 98.9%, while personnel competency improved from 68.0% to 88.0%. Knowledge retention increased from 71.0% to 84.0%, indicating improved sustained understanding of documentation procedures. The composite GMP compliance index increased from 76.0 to 90.0 points. These illustrative outcomes indicate how DERTOA-guided training could improve documentation performance through targeted competency development, prioritization of high-risk activities, and systematic monitoring of recurrent deficiencies. The proposed evaluation distinguishes immediate training gains from sustained operational improvements using longitudinal measurements and comparative statistical analysis.

Table 4 Simulated Pre-Training and Post-Training Documentation Quality, Personnel Competency, Knowledge Retention, and GMP Compliance Indicators

Performance Indicator	Pre-Training	Post-Training	Improvement
Documentation error rate (per 1,000 opportunities)	25.0	11.0	56.0% reduction
Documentation accuracy (%)	97.5	98.9	+1.4 pp
Personnel competency score (%)	68.0	88.0	+20.0 pp
Knowledge retention at 90 days (%)	71.0	84.0	+13.0 pp
GMP compliance index (0–100)	76.0	90.0	+14.0 points




Fig 5 Simulated Documentation Error Reduction (%) Under Five Algorithm-Guided Training Allocation Strategies Across Four Post-Training Follow-Up Periods.

Figure 5 compares simulated documentation error reductions achieved through five training allocation strategies across 30, 60, 90, and 180 days. DERTOA demonstrates the highest illustrative reduction, declining from 56% at 30 days to 53%, 50%, and 46% during subsequent assessments. XGBoost-guided allocation records corresponding reductions of 44%, 42%, 39%, and 35%, while Random Forest achieves 36%, 34%, 32%, and 28%. Logistic Regression produces reductions of 28%, 26%, 24%, and 20%, compared with 20%, 18%, 16%, and 12% under conventional rule-based allocation. DERTOA maintains a 34-percentage-point advantage over the rule-based strategy at 180 days. Its mean reduction across follow-up periods is 51.25%, compared with 40.00% for XGBoost and 16.50% for rule-based allocation. The simulated decline over time illustrates the importance of competency retention monitoring, refresher interventions, and adaptive training optimization for sustaining documentation improvements.

- *Comparative Predictive Performance of DERTOA Against Logistic Regression, Random Forest, XGBoost, and Conventional Rule-Based Approaches Using Classification and Regression Performance Metrics*

The comparative predictive assessment evaluates DERTOA against four established algorithms using standardized classification and regression performance indicators. Table 4.3 presents illustrative simulation results indicating that DERTOA achieves the highest F1-score of 93.1%, an AUC-ROC of 97.0%, and the lowest mean absolute error (MAE) of 1.8 errors per 1,000 documentation opportunities. XGBoost follows with an F1-score of 86.0% and MAE of 2.6, while Random Forest records 80.7% and 3.5, respectively. Logistic Regression achieves an F1-score of 72.2%, compared with 62.5% for rule-based detection. These differences illustrate the proposed benefits of integrating severity-weighted documentation risk features, nonlinear prediction, and personnel competency indicators. The comparative framework also evaluates prediction calibration and error magnitude to distinguish classification effectiveness from quantitative documentation error forecasting accuracy under identical testing conditions.

Table 5 Simulated Comparative Classification and Regression Performance of DERTOA and Benchmark Algorithms

Algorithm	F1-Score (%)	AUC-ROC (%)	MAE (Errors/1,000)
Conventional Rule-Based	62.5	68.0	5.8
Logistic Regression	72.2	77.0	4.7
Random Forest	80.7	85.0	3.5
XGBoost	86.0	91.0	2.6
DERTOA	93.1	97.0	1.8




Fig 6 Simulated Comparative Predictive Performance of Five Documentation Error Assessment Algorithms Using Precision, Recall, F1-score, AUC-ROC, and Normalized Regression Accuracy.

Figure 6 illustrates the simulated predictive performance of five algorithms across classification and regression indicators. DERTOA achieves the highest precision (94.0%), recall (92.3%), F1-score (93.1%), and AUC-ROC (97.0%), indicating stronger illustrative discrimination between compliant and non-compliant documentation records. XGBoost follows with precision of 87.0%, recall of 85.0%, F1-score of 86.0%, and AUC-ROC of 91.0%. Random Forest records 82.0% precision and 79.5% recall, while Logistic Regression achieves 74.0% and 70.5%, respectively. Conventional rule-based detection demonstrates the lowest recall at 61.1%. DERTOA also records the lowest MAE of 1.8 errors per 1,000 opportunities, compared with 2.6 for XGBoost and 5.8 for rule-based detection. This represents a 30.8% lower MAE than XGBoost. The radar chart therefore illustrates the hypothesized advantage of integrated documentation risk modeling, although empirical validation is required to establish reproducibility and statistical significance.

- *Adaptive Training Optimization, Compliance Risk Reduction, Statistical Significance, Model Interpretability, Sensitivity Analysis, and Implications for Sustainable GMP Compliance*

The simulated assessment evaluates adaptive training optimization and sustained documentation risk reduction across five computational approaches. Table 6 indicates that DERTOA achieved the greatest initial documentation error reduction of 56.0%, compared with 44.0% for XGBoost, 36.0% for Random Forest, 28.0% for Logistic Regression, and 20.0% for conventional rule-based allocation. After 180 days, DERTOA retained a 46.0% error reduction, demonstrating the strongest illustrative sustained performance. Its predictive recall of 92.3% further supports the proposed integration of severity-weighted risk assessment, personnel competency monitoring, and adaptive training prioritization. Model interpretability is supported through documented risk contributors and auditable training recommendations. Statistical significance and sensitivity to training budgets, risk weights, and operational workloads require independent testing. These comparisons establish a quantitative framework for evaluating whether targeted interventions can sustain GMP documentation improvements while reducing compliance risks and training resource inefficiencies.

Table 6 Comparative Simulated Adaptive Training Optimization, Sustained Documentation Error Reduction, and Predictive Recall

Algorithm	Error Reduction at 30 days (%)	Error Reduction at 180 days (%)	Predictive Recall (%)
Conventional Rule-Based	20.0	12.0	61.1
Logistic Regression	28.0	20.0	70.5
Random Forest	36.0	28.0	79.5

XGBoost	44.0	35.0	85.0
DERTOA	56.0	46.0	92.3

Figure 7 presents a comparative heatmap of five algorithms across documentation error reduction and predictive performance indicators. DERTOA records the strongest illustrative results, achieving 56.0% error reduction at 30 days and maintaining 46.0% at 180 days. Its predictive recall, F1-score, and AUC-ROC reach 92.3%, 93.1%, and 97.0%, respectively. XGBoost achieves corresponding error reductions of 44.0% and 35.0%, with recall of 85.0%, F1-score of 86.0%, and AUC-ROC of 91.0%. Random Forest produces 36.0% initial error reduction and 28.0% sustained reduction, whereas Logistic Regression records 28.0% and 20.0%. Rule-based allocation demonstrates the lowest reductions of 20.0% and 12.0%. DERTOA therefore maintains an illustrative 11-percentage-point advantage over XGBoost in sustained error reduction. These patterns suggest potential benefits from adaptive training allocation, severity-weighted risk prediction, and competency monitoring, although statistical significance and operational effectiveness remain unverified.




Fig 7 Comparative Simulated Documentation Error Reduction, Sustained Training Effectiveness, and Predictive Performance of Five Algorithms Across GMP Compliance Indicators.

- *Statistical Significance, Interpretability, and Sensitivity Analysis*

The quantitative comparison also provides a basis for examining whether DERTOA's apparent advantages remain stable under changing training resources and documentation workloads.

Table 7 Statistical Significance, Interpretability, and Sensitivity Analysis

Evaluation Dimension	Proposed Assessment	Interpretation Criterion
Statistical significance	Cluster-aware confidence intervals and paired model comparisons	Determine whether differences are distinguishable from sampling variability
Model interpretability	Feature attribution and documented risk scores	Identify factors contributing to predictions and training recommendations
Budget sensitivity	Training budget varied by ±20%	Evaluate allocation stability and compliance risk trade-offs
Risk-weight sensitivity	Severity weights varied by ±20%	Assess robustness of documentation risk priorities
Operational sustainability	Repeated evaluation at 30, 60, 90, and 180 days	Assess persistence of training-related improvements

These analyses are proposed validation procedures, not completed statistical tests. In particular, a statistically significant training effect cannot be inferred from the five algorithm-level summary values alone.

V. CONCLUSIONS AND RECOMMENDATIONS

- *Summary of Principal Findings on Laboratory Documentation Errors, Personnel Training Effectiveness, and Quantitative GMP Compliance Performance*

The study developed a quantitative framework for examining laboratory documentation deficiencies, personnel training effectiveness, and Good Manufacturing Practice (GMP) compliance within pharmaceutical quality control operations. The illustrative simulation identified substantial differences in documentation error frequency, severity, and operational distribution across laboratory activities. Sample registration exhibited the highest documentation error rate of 24.6 errors per 1,000 opportunities, while analytical testing recorded the highest proportion of high-severity deficiencies at 31.4%. These patterns demonstrated the importance of evaluating documentation performance through both error frequency and potential regulatory consequences. The simulated longitudinal training assessment indicated that documentation error rates decreased from 25.0 to 11.0 errors per 1,000 opportunities, representing a 56.0% reduction. Personnel competency improved from 68.0% to 88.0%, knowledge retention increased from 71.0% to 84.0%, and the composite GMP compliance index increased from 76.0 to 90.0.

Comparative algorithm evaluation demonstrated the proposed capabilities of the Documentation Error Risk and Training Optimization Algorithm (DERTOA). Under the illustrative simulation, DERTOA achieved precision of 94.0%, recall of 92.3%, F1-score of 93.1%, AUC-ROC of 97.0%, and mean absolute error of 1.8 errors per 1,000 documentation opportunities. Its simulated 180-day documentation error reduction remained at 46.0%, compared with 35.0% for XGBoost and 12.0% for conventional rule-based allocation. These findings illustrate the potential advantages of integrating predictive risk assessment, severity-weighted documentation indicators, competency retention monitoring, and adaptive training optimization. Nevertheless, the reported numerical outcomes represent synthetic demonstrations rather than independently validated pharmaceutical laboratory results, and their practical significance requires prospective evaluation.

- *Contributions of the Proposed DERTOA Algorithm to Documentation Error Prediction, Risk-Based Training Optimization, and Pharmaceutical Quality Assurance*

The principal methodological contribution of this research is the proposed Documentation Error Risk and Training Optimization Algorithm (DERTOA), which integrates predictive analytics, severity-weighted documentation risk assessment, personnel competency evaluation, and constrained multi-objective training optimization into a unified pharmaceutical quality assurance framework. Unlike conventional compliance assessment systems that predominantly identify deficiencies through retrospective inspections and predefined procedural thresholds, DERTOA is designed to estimate documentation non-compliance probabilities and prioritize interventions according to operational risk. Its architecture combines gradient-boosted prediction with documentation severity scores, historical recurrence indicators, competency retention measurements, and training resource constraints. This integrated structure enables risk-based allocation of corrective training while maintaining transparent documentation of the factors influencing intervention recommendations.

The simulated comparative evaluation illustrated DERTOA's potential advantages over Logistic Regression, Random Forest, XGBoost, and rule-based detection. Its F1-score of 93.1% exceeded the corresponding values of 86.0%, 80.7%, 72.2%, and 62.5%, respectively. DERTOA also produced an illustrative 30.8% reduction in mean absolute error relative to XGBoost. Beyond predictive classification, the proposed algorithm introduces a training optimization mechanism intended to balance residual documentation risk, instructional costs, employee competency deficiencies, and operational availability. Its anticipated contribution to pharmaceutical quality assurance involves earlier identification of documentation vulnerabilities, improved prioritization of corrective and preventive actions, and continuous monitoring of workforce competency. The framework further supports interpretable risk reporting through feature attribution, documented intervention priorities, and compliance performance indicators. These contributions establish a testable computational methodology for intelligent GMP documentation management, although claims of practical superiority require independent validation using representative laboratory records and controlled training interventions.

- *Recommendations for Risk-Based Documentation Control, Continuous Personnel Competency Assessment, Training Program Improvement, and Regulatory Inspection Readiness*

Pharmaceutical manufacturers should establish integrated documentation quality monitoring systems that combine standardized error classification, severity-based risk assessment, systematic personnel competency evaluation, and continuous GMP compliance surveillance. Laboratory documentation audits should prioritize activities exhibiting elevated error frequencies, recurring deficiencies, or substantial potential consequences for analytical validity and product quality decisions. Particular attention should be directed toward sample identification, analytical testing records, result transcription, instrument calibration documentation, and electronic audit-trail verification. Documentation performance should be measured using standardized error rates per 1,000 audited opportunities, severity-weighted risk indices, correction timeliness, and recurrence indicators. These measurements should be incorporated into laboratory quality management dashboards to support consistent departmental comparisons and early identification of emerging compliance vulnerabilities.

Training programs should progress beyond attendance-based evaluation toward demonstrable

competency improvement and sustained documentation accuracy. Personnel assessments should incorporate baseline examinations, supervised practical demonstrations, immediate post-training evaluations, and delayed knowledge retention measurements. Where operationally appropriate, DERTOA-guided training recommendations could prioritize employees and analytical activities according to validated documentation risk indicators and identified competency gaps. Quality assurance departments should retain responsibility for approving training assignments, reviewing predictive outputs, and determining corrective and preventive actions. Manufacturers should additionally strengthen electronic record access controls, audit-trail review procedures, version-controlled documentation systems, and periodic data integrity assessments. Regulatory inspection readiness should be supported by accessible training histories, complete analytical records, documented deviation investigations, approved corrective actions, and evidence of sustained compliance monitoring. Pilot implementation should begin within selected laboratory activities, followed by controlled performance evaluation before broader operational adoption.

- *Study Limitations, Model Generalizability, Future Research Directions, and Opportunities for Intelligent GMP Compliance Monitoring*

Despite its methodological contributions, the proposed research framework has limitations concerning data availability, simulation assumptions, predictive model generalizability, and the causal interpretation of training effectiveness. The numerical results presented in the study were developed as illustrative synthetic comparisons and therefore cannot establish actual documentation error prevalence, algorithmic superiority, or training-related compliance improvements in pharmaceutical manufacturing facilities. Synthetic datasets may not adequately reproduce the complexity of genuine laboratory operations, including rare critical documentation violations, inconsistent audit practices, changing analytical workloads, heterogeneous personnel competencies, and regulatory inspection variability. Furthermore, predictive models trained on records from individual laboratories may perform differently across facilities employing alternative documentation procedures, electronic quality management systems, or workforce training arrangements. The proposed severity-weighted compliance index also depends on predefined scoring criteria, requiring expert validation and sensitivity analysis before practical implementation.

Future research should prioritize multicentre longitudinal studies incorporating authentic GMP laboratory documentation records, independent external validation, and prospectively evaluated training interventions. Comparative investigations should examine DERTOA performance under varying laboratory workloads, documentation complexity, training budgets, personnel experience levels, and regulatory risk classifications. Additional methodological developments could incorporate explainable machine learning, temporal anomaly detection, competency decay modeling, uncertainty quantification, and human-in-the-loop decision support. Controlled or staggered intervention studies would help distinguish genuine training effects from concurrent operational changes. Integration with laboratory information management systems and electronic quality management platforms could facilitate automated monitoring of documentation completeness, correction delays, and recurring deficiencies. Ultimately, the development of intelligent GMP compliance monitoring should emphasize predictive reliability, operational transparency, regulatory accountability, and demonstrable improvements in pharmaceutical documentation quality rather than predictive performance alone.

REFERENCES

[1]. Amebleh, J., Igba, E., & Ijiga, O. M. (2021). Graph-based fraud detection in open-loop gift cards: Heterogeneous GNNs, streaming feature stores, and near-zero-lag anomaly alerts. International Journal of Scientific Research in Science, Engineering and Technology, 8(6), 317–339. https://doi.org/10.32628/IJSRSET214418

[2]. Bell, B. S., Tannenbaum, S. I., Ford, J. K., Noe, R. A., & Kraiger, K. (2017). 100 years of training and development research: What we know and where we should go. Journal of Applied Psychology, 102(3), 305–323. https://doi.org/10.1037/apl0000142

[3]. Biau, G., & Scornet, E. (2016). A random forest guided tour. TEST, 25(2), 197–227. https://doi.org/10.1007/s11749-016-0481-7

[4]. Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (pp. 785–794). Association for Computing Machinery. https://doi.org/10.1145/2939672.2939785

[5]. Gama, J., Žliobaitė, I., Bifet, A., Pechenizkiy, M., & Bouchachia, A. (2014). A survey on concept drift adaptation. ACM Computing Surveys, 46(4), Article 44. https://doi.org/10.1145/2523813

[6]. Grangeia, H. B., Silva, C., Simões, S. P., & Reis, M. S. (2020). Quality by design in pharmaceutical manufacturing: A systematic review of current status, challenges and future perspectives. European Journal of Pharmaceutics and Biopharmaceutics, 147, 19–37. https://doi.org/10.1016/j.ejpb.2019.12.007

[7]. Hughes, A. M., Zajac, S., Spencer, J. M., & Salas, E. (2018). A checklist for facilitating training transfer in organizations. International Journal of Training and Development, 22(4), 334–345. https://doi.org/10.1111/ijtd.12141

[8]. Hughes, A. M., Zajac, S., Woods, A. L., & Salas, E. (2020). The role of work environment in training sustainment: A meta-analysis. Human Factors, 62(1), 166–183. https://doi.org/10.1177/0018720819845988

[9]. Idika, C. N., Salami, E. O., Ijiga, O. M., & Enyejo, L. A. (2021). Deep learning driven malware classification for cloud-native microservices in edge computing architectures. International Journal of Scientific Research in Computer Science, Engineering and Information Technology, 7(4). https://doi.org/10.32628/CSEIT182551

[10]. Ijiga, O. M., Ifenatuora, G. P., & Olateju, M. (2021a). Bridging STEM and cross-cultural education: Designing inclusive pedagogies for multilingual classrooms in Sub-Saharan Africa. IRE Journals, 5(1), 543–559.

[11]. Ijiga, O. M., Ifenatuora, G. P., & Olateju, M. (2021b). Digital storytelling as a tool for enhancing STEM engagement: A multimedia approach to science communication in K-12 education. International Journal of Multidisciplinary Research and Growth Evaluation, 2(5), 495–505. https://doi.org/10.54660/.IJMRGE.2021.2.5.495-505

[12]. Ijiga, O. M., Ifenatuora, G. P., & Olateju, M. (2022). AI-powered e-learning platforms for STEM education: Evaluating effectiveness in low-bandwidth and remote learning environments. International Journal of Scientific Research in Computer Science, Engineering and Information Technology, 8(5), 455–475. https://doi.org/10.32628/CSEIT23902187

[13]. Jaiswal, H., Muddukrishna, B. S., & Kulyadi, G. P. (2020). Data integrity violations: A challenge to the pharmaceutical industry. International Journal of Pharmaceutical Quality Assurance, 11(1), 196–198. https://doi.org/10.25258/ijpqa.11.1.30

[14]. James, U. U. (2022). Federated identity management using secure enclaves for cross-domain access control in 5G edge networks. International Journal of Scientific Research in Science, Engineering and Technology, 9(6), 450–471. https://doi.org/10.32628/IJSRSET25122271

[15]. Kwarteng, R. A., Idoko, I. P., Ijiga, O. M., & Enyejo, L. A. (2020). Integrating cybersecurity awareness and access control into organizational IT operations for risk reduction. International Journal of Scientific Research in Computer Science, Engineering and Information Technology, 6(1), 243–261. https://doi.org/10.32628/CSEIT23906128

[16]. Lacerenza, C. N., Reyes, D. L., Marlow, S. L., Joseph, D. L., & Salas, E. (2017). Leadership training design, delivery, and implementation: A meta-analysis. Journal of Applied Psychology, 102(12), 1686–1718. https://doi.org/10.1037/apl0000241

[17]. Mrazek, C., Lippi, G., Keppel, M. H., Felder, T. K., Oberkofler, H., Haschke-Becher, E., & Cadamuro, J. (2020). Errors within the total laboratory testing process, from test selection to medical decision-making: A review of causes, consequences, surveillance and solutions. Biochemia Medica, 30(2), Article 020502. https://doi.org/10.11613/BM.2020.020502

[18]. Nwokocha, C. R., Peter-Anyebe, A. C., & Ijiga, O. M. (2021). Optimizing agile-based system integration for enhanced ECMS functionality and Smile CDR adoption within health information networks. International Journal of Scientific Research in Computer Science, Engineering and Information Technology, 7(6), 470–490. https://doi.org/10.32628/CSEIT2282148

[19]. Ogundola, T. (2020). Advances in quantum-secure banking: Cryptographic solutions. International Journal of Science and Research Archive, 1(1), 228–230. https://doi.org/10.30574/ijsra.2020.1.1.0048

[20]. Ogundola, T. (2022). Quantum-resistant cryptographic techniques for financial institutions. International Journal of Scientific Research and Modern Technology, 1(11), 18–20. https://doi.org/10.38124/ijsrmt.v1i11.586

[21]. Onyekaonwu, C. B., & Peter-Anyebe, A. C. (2019). Empowering underserved youth through tech-enabled STEM education: A participatory model from Nigeria. International Journal of Scientific Research in Science and Technology, 6(5). https://doi.org/10.32628/IJSRST1965990

[22]. Onyekaonwu, C. B., Peter-Anyebe, A. C., & Raphael, F. O. (2019). From prescription to prediction: Leveraging AI/ML to improve medication adherence and adverse drug event detection in community pharmacies. International Journal of Scientific Research in Science and Technology, 6(6), 460–476. https://doi.org/10.32628/IJSRST52310387

[23]. Plebani, M. (2015). Diagnostic errors and laboratory medicine: Causes and strategies. EJIFCC, 26(1), 7–14.

[24]. Plebani, M., Sciacovelli, L., & Aita, A. (2017). Quality indicators for the total testing process. Clinics in Laboratory Medicine, 37(1), 187–205. https://doi.org/10.1016/j.cll.2016.09.015

[25]. Rantanen, J., & Khinast, J. (2015). The future of pharmaceutical manufacturing sciences. Journal of Pharmaceutical Sciences, 104(11), 3612–3638. https://doi.org/10.1002/jps.24594

[26]. Rogers, C. A., Ahearn, J. D., & Bartlett, M. G. (2020). Data integrity in the pharmaceutical industry: Analysis of inspections and warning letters issued by the Bioresearch Monitoring Program between fiscal years 2007–2018. Therapeutic Innovation & Regulatory Science, 54(5), 1123–1133. https://doi.org/10.1007/s43441-020-00129-z

[27]. Rudin, C. (2019). Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. Nature Machine Intelligence, 1, 206–215. https://doi.org/10.1038/s42256-019-0048-x

[28]. Saito, T., & Rehmsmeier, M. (2015). The precision-recall plot is more informative than the ROC plot when evaluating binary classifiers on imbalanced datasets. PLOS ONE, 10(3), Article e0118432. https://doi.org/10.1371/journal.pone.0118432

[29]. Schmitt, J., Bönig, J., Borggräfe, T., Beitinger, G., & Deuse, J. (2020). Predictive model-based quality

inspection using machine learning and edge cloud computing. Advanced Engineering Informatics, 45, Article 101101. https://doi.org/10.1016/j.aei.2020.101101

[30]. Sciacovelli, L., Lippi, G., Sumarac, Z., Garcia Del Pino Castro, I., Ivanov, A., De Guire, V., Coskun, C., Aita, A., Padoan, A., & Plebani, M. (2019). Pre-analytical quality indicators in laboratory medicine: Performance of laboratories participating in the IFCC working group "Laboratory Errors and Patient Safety" project. Clinica Chimica Acta, 497, 35–40. https://doi.org/10.1016/j.cca.2019.07.007

[31]. Sciacovelli, L., Lippi, G., Sumarac, Z., West, J., Garcia Del Pino Castro, I., Furtado Vieira, K., Ivanov, A., & Plebani, M. (2017). Quality indicators in laboratory medicine: The status of the progress of IFCC Working Group "Laboratory Errors and Patient Safety" project. Clinical Chemistry and Laboratory Medicine, 55(3), 348–357. https://doi.org/10.1515/cclm-2016-0929

[32]. U.S. Food and Drug Administration. (2018). *Data integrity and compliance with drug CGMP: Questions and answers. Guidance for industry.* https://www.fda.gov/regulatory-information/search-fda-guidance-documents/data-integrity-and-compliance-drug-cgmp-questions-and-answers n

[33]. West, J., Atherton, J., Costelloe, S. J., Pourmahram, G., Stretton, A., & Cornes, M. (2017). Preanalytical errors in medical laboratories: A review of the available methodologies of data collection and analysis. Annals of Clinical Biochemistry, 54(1), 14–19. https://doi.org/10.1177/0004563216669384

[34]. Yu, L. X., Amidon, G., Khan, M. A., Hoag, S. W., Polli, J., Raju, G. K., & Woodcock, J. (2014). Understanding pharmaceutical quality by design. The AAPS Journal, 16(4), 771–783. https://doi.org/10.1208/s12248-014-9598-3